

## Suboptimal Healthcare for the Homeless

High rates of leaving against medical advice



9.3% of homeless patients leave the hospital AMA, compared with the national average of 1.3%.



Homelessness is one of the greatest predictors of Emergency Department (ED) super-utilization



Homeless persons are 3x more likely than members of the general public to use ED services ≥1x per year.

ED Use:
3X
greater odds

**30-day Hospital Readmission** 





Average Cost \$13,800

Within 30 days of leaving the hospital, >20% of homeless patients were readmitted, compared to 7% among matched controls.

## Stigma and Judgment

"I didn't feel like a ... person, I just felt like I was part of their job and they shoved me through. Get me in and out as fast as they can. A kind of process."

(Nickasch and Marnocha, 2009)

## References

- 1) Barrett ML, Wier LM, Jiang HJ, et al. (2015). "All-Cause Readmissions by Payer and Age, 2009–2013: Statistical Brief #199." Healthcare Cost and Utilization Project (HCUP) Statistical Briefs [Internet]. Agency for Healthcare Research and Quality (US)
- 2) Gindi RM, Black LI, Cohen RA. (2016). "Reasons for emergency room use among U.S. adults aged 18–64: National Health Interview Survey, 2013 and 2014." *National health statistics reports; no 90.*
- 3) n.a. (16 Nov. 2018). "NHS Not Tracking Why More Patients Are Being Readmitted to Hospital." *Healthwatch Central Bedfordshire*.
- 4) Nickasch, Bonnie, and Suzanne K. Marnocha. (2009). "Healthcare Experiences of the Homeless." *Journal of the American Academy of Nurse Practitioners.*
- 5) Saab, Dima, et al. (2016). "Hospital Readmissions in a Community-Based Sample of Homeless Adults: a Matched-Cohort Study." *Journal of General Internal Medicine*.
- 6) Sun, Ruirui, et al. (2017). "Characteristics of Homeless Individuals Using Emergency Department Services in 2014." AHRQ.